



AFTER CARE



AFTER CARE



AFTER CARE

Thank you for choosing REVIV as your wellness provider. As a quick reminder, even though the vast majority of patients receiving IV or Booster Shot therapies feel greatly improved afterwards, every individual is unique and there is no guarantee that you will notice a difference in the way you feel after a therapy, nor does improvement of symptoms exclude other coexisting potential medical conditions.

You have just received an intravenous (IV) and or an intramuscular (IM) therapy. Your therapy may consist of vitamins, anti-oxidants, minerals, electrolytes, amino-acids, fluids and sometimes medications or any combination of the above.

Home Care Instructions

- Leave your bandage on for at least 15 minutes after your therapy
- Change your bandage if there is a small amount of bleeding at the puncture site
- Keep the puncture site area clean and dry after you remove your bandage
- Avoid showering, bathing or exposing the puncture site to water for at least 1 hour after your therapy
- Avoid strenuous activity or stressing the puncture site for 24 hours after your therapy

Procedure Risks and Complications

- Pain at the puncture site
- Bruising of the skin
- Infection
- Allergic reaction to any of the ingredients from the procedure preparation or the therapy
- Failure to successfully place IV catheter in the vein
- Puncture of the wall of the vein requiring the use of a different site for IV
- Self-resolving blood clot in the superficial vein

- Infiltration – fluid and ingredient leakage through the wall of a vein into the surrounding skin, which will reabsorb spontaneously

Contact Us or Seek Medical Care

- If you develop any loss of sensation or numbness around the procedure area or in the arms where the procedure was performed
- If you develop repeated or worsening swelling at the puncture site
- If there is bleeding from the puncture area that will not stop
- If a large bruise develops around the puncture site
- If bleeding from the puncture site develops and will not stop even after pressure is applied for 15 minutes
- If redness or red streaking develops around the procedure site
- If you develop shortness of breath, rash, itching sensation, swelling of your lips, tongue, throat or other signs of an allergic reaction
- If you develop a fever

Contact: care@revivme.com

