




— · VACCINATION RECORD —



PERSONAL DETAILS

Name

Date of Birth

Home Address

Phone Number

Known Allergies

Blood Type & Medications

TRAVEL VACCINATIONS



Vaccination	Date Received	Brand	Given By (Signature)	Booster/ Re-Vaccination
DIPHTHERIA, TETANUS & POLIO				
TYPHOID				
HEPATITIS A				
Dose 1				
Booster				

TRAVEL VACCINATIONS



Vaccination	Date Received	Brand	Given By (Signature)	Booster/ Re-Vaccination
-------------	---------------	-------	----------------------	-------------------------

HEPATITIS B

Dose 1				
Dose 2				
Dose 3				

CHOLERA

Dose 1				
Dose 2				
Dose 1				
Dose 2				

Vaccination	Date Received	Brand	Given By (Signature)	Booster/ Re-Vaccination
-------------	---------------	-------	----------------------	-------------------------

RABIES

Dose 1				
Dose 2				
Dose 3				
Dose 4				

JAPANESE ENCEPHALITIS

Dose 1				
Dose 2				
Dose 1				
Dose 2				

TRAVEL VACCINATIONS

GENERAL VACCINATIONS



Brand	Date Received	Given By (Signature)	Dosing Schedule	Total No. of Tablets
MALARIA				
Malarone			Once daily for two days before entering an at risk area, then for the duration of stay within that area and for seven days after leaving/ returning home	
Malarone			Once daily for two days before entering an at risk area, then for the duration of stay within that area and for seven days after leaving/ returning home	
Malarone			Once daily for two days before entering an at risk area, then for the duration of stay within that area and for seven days after leaving/ returning home	
Malarone			Once daily for two days before entering an at risk area, then for the duration of stay within that area and for seven days after leaving/ returning home	

Vaccination	Date Received	Brand	Given By (Signature)	Booster/ Re-Vaccination
HPV				
Dose 1				
Dose 2				
Dose 3				
FLU				
PNEUMONIA (POLYSACCHARIDE)				
PNEUMONIA (CONJUGATE)				



AFTERCARE ADVICE

Like any medications, vaccines can cause side effects.

Sometimes this can include redness and tenderness at the injection site, and less commonly, they can cause slight fever, malaise, headache and generalised aching. These side effects usually last 12-24 hours and fever and aches can be reduced with paracetamol and plenty of non-alcoholic fluids. Please seek medical advice if you are concerned, either from your GP or call REVIV.

If you have side effects please report them via the Yellow Card Scheme at www.yellowcard.mhra.gov.uk

CONTACT

care@revivme.com

VISIT

revivme.com @ f [revivme](#) [revivuk](#)

IV THERAPY • BOOSTER SHOTS • LIFESTYLE GENETICS
VACCINATIONS • BLOOD TESTING

